

TO ALL TO WHOM THESE PRESENTS SHAVE COME;

Airginin Çech Intellectual Properties, Inc.

THETERS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT. SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS ARROW SELUNG THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT,

CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN UCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY CITON ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CRITIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF A STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

BARLEY

'Price

In Testimonn Mercest, I have hereunto set my hand and caused the seal of the Mant Mariety Frotestion Office to be affixed at the City of Washington, D.C. this sixteenth day of September, in the year two thousand three.

Rangele

Commissioner Plant Varioty Protection Office Apricultural Marketing Service

riculturo

Z0030013Z

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,705 (\$320 filing fee and \$2,385 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initiated and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$320 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM

18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Foundation seed of Price barley was first provided to seedsmen for multiplication in Fall 2002.

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political baliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

ST_470 (10.00) designed by the Plant Variety Projection Office with Word 2000. Replaces STD_470 (07.01) which is obsolete



'PRICE' WINTER FEED BARLEY

18A. EXHIBIT A: ORIGIN AND BREEDING HISTORY

Price winter feed barley was derived from the cross 'Callao'/SC830366. The experimental line SC830366, derived from the cross 'McNair 601'/'Harrison'// 'Gembloux'/3/'Henry' and developed at Clemson University, was selected as a parent from the 1988-89 USDA-ARS Uniform Winter Barley Yield Nursery. The cross was made in spring1990, and the F₁ generation was grown in the field as a single 4ft headrow in 1991 to produce F₂ seed. The population was advanced from the F₂ to F₄ generation using a modified bulk breeding method.

POPULATION ADVANCEMENT AND SELECTION OF THE VARIETY

Barley spikes were selected from the population in each segregating generation (F_2-F_4) on the basis of absence of obvious disease, early maturity, short straw and desirable head type and size. Selected spikes were threshed in bulk, and the seed was planted in $225 \, \mathrm{ft}^2$ blocks in the fall of each year. Spikes selected from the F_4 bulk where threshed individually and planted in separate 4ft headrows. Price was derived as a bulk of one of these F_5 headrows selected in 1995. The line was tested as entry 321 in non-replicated observation tests in 1996 and was designated VA96-44-321. This line was tested in Virginia's Official Variety Trials from 1997 to 2001 (Tables 1-6). It also was tested for three years (1998-2000) in the USDA-ARS Uniform Winter Barley Yield Nursery (Tables 7-12) and in the USDA-ARS Uniform Barley Winter Hardiness Nursery (Tables 13-15).

MULTIPLICATION AND PURIFICATION

Initial Breeder Seed of Price was developed in 2000-01 via removal of variant plants from a 0.25 acre F₁₁ increase block planted at the VCIA Foundation Seed Farm. This increase block produced 30 bu of Breeder Seed, of which 22 bu were planted on 10 acres at VCIA Foundation Seed Farm in fall 2001. The field was scouted and variant plants were removed during the growing season. The field was harvested and produced 1100 bu of Price Foundation Seed. A portion of this seed was planted on 16 acres at the Foundation Seed Farm in fall 2002, and the remaining seed was provided to seedsmen for multiplication. While Price has remained stable and uniform in composition through the last two generations of multiplication, the initial Breeder Seed contained the following proportion of variants: up to 0.1% taller plants, 0.1% plants having no awns or short awnlets, and 0.1% plants having long awns.

Development of a purer source of Price Breeder Seed was initiated in fall 2002. During fall 2002, 400 headrows of Price, each originating from a single spike, were planted and will be evaluated in 2003 for purity and trueness of type. Variant headrows will be removed prior to harvest, and the remaining rows will be harvested and advanced to form a purer source of Breeder seed.

'PRICE' WINTER FEED BARLEY

18B. EXHIBIT B: NOVELTY STATEMENT

Price is uniquely different from all know barley cultivars, but is most similar to its parent Callao. On the basis of disease reaction (0=immune to 4=highly susceptible) of seedlings in greenhouse tests conducted from 1997-2000, Price is highly resistant (infection type = 0;N) to leaf rust (*Puccinia hordei*) race 8, while Callao is moderately susceptible (IT=23). Conversely, seedlings of Price are moderately susceptible (IT=3,4) to powdery mildew (*Blumeria graminis* f. sp. hordei), while those of Callao are highly resistant (IT=0). On average, head emergence of Price is 3.3 days (range 2-5 days) later than Callao (See table below). Average plant height of Price is 1.9 inches (range 1-3 inches) taller than Callao. Straw strength of Price is better than that of Callao on the basis of lodging scores (0.2=no lodging to 10=complete lodging). Average lodging score of Price (1.4) was lower than that of Callao (5.9) with differences ranging from 3.4 to 5.6 over the five year period.

Price versus Callao for Heading Date, Plant Height and Lodging

	Price	Callao	Difference	L.S.D. (0.05)
5-Year Mean(1997-2001)	:			
Heading Date (Mar 31+)	20.6	17.3	3.3*	0.4
Plant Height (inches)	35.6	33.7	1.9*	0.5
Lodging (0.2 – 10)	1.4	5.9	4.5*	0.5
1996-1997 Season				
Heading Date (Mar 31+)	23	20	3*	1.0
Plant Height (inches)	37	36	1*	1.0
Lodging (0.2 – 10)	1.2	5.0	3.8*	1.1
1997-1998 Season				
Heading Date (Mar 31+)	18	15	3*	1.0
Plant Height (inches)	36	33	3*	1.0
Lodging (0.2 – 10)	2.8	7.6	4.8*	1.2
1998-1999 Season				
Heading Date (Mar 31+)	22	19	3*	1.0
Plant Height (inches)	37	35	2*	1.0
Lodging (0.2 – 10)	0.3	4.0	3.7*	0.6
1999-2000 Season				
Heading Date (Mar 31+)	17	15	2*	1.0
Plant Height (inches)	37	35	2*	1.0
Lodging (0.2 – 10)	1.0	4.4	3.4*	0.8
2000-2001 Season				
Heading Date (Mar 31+)	25	20	5*	1.0
Plant Height (inches)	31	30	1*	1.0
Lodging (0.2 – 10)	2.1	7.7	5.6*	1.1

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK AND SEED DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C (Barley)

OBJECTIVE DESCRIPTION OF VARIETY

INSTRUCTIONS: Joe Reverse,	LOW FOLONIE!
Virginia Tech Intellectual Properties, Inc.	FOR OFFICIAL USE ONLY
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	700300132
1872 Pratt Drive, Suite 1625	VARIETY NAME OR TEMPORARY
Blacksburg, VA 24060	OESIGNATION Price
Place the appropriate number that describes the varietal character	
Place a zero in first box (i.e. 0 8 9 or 0 9) when number i	s either 99 or less or 9 or less.
1. GROWTH HABIT:	
3 1 - SPRING 2 - FACULTATIVE WINTER 3 - WINTER	2 Early Growth: 1 - PROSTRATE 2 - SEMIPROSTRATE 3 - ERECT
2. MATURITY (50% Flowering):	
2 1 = EARLY (California Mariout) 2 = MIDSEASON (Betzes)	3 = LATE (Frontier)
No. of days Earlier than 8 1 - BETZES 2 - CAL	IFORNIA MARIOUT 3 = CONQUEST 4 = DICKSON
No. of days Later than 9 5 - PIROLINE 6 - PR	MMUS 7-UNITAN 8=Wysor 9=Callao
3, PLANT HEIGHT (From soil level to top of head):	
2 1 - SEMIDWARF 2 - SHORT (Cellfornia Meriout) 3 - MED	IUM TALL (Betzes) 4 = TALL (Conquest)
(LIFORNIA MARIOUT 3 - CONQUEST 4 - DICKSON RIMUS 7 - UNITAN 8=Wysor 9=Callao
5 Cm. Taller than 9	
4. STEM:	
2 Exertion (Flue to spike at maturity): 3 = 10 - 15 cm.	1 Anthocyanin: 1 - ABSENT 2 - PRESENT
0 5 NO, OF NODES (Originating from node above ground).	
1 - CLOSED 2 - V-SHAPED 3 - OPEN 1 - CLOSED 2 - V-SHAPED 3 - OPEN 4 - MODIFIED CLOSED OR OPEN	1 - STRAIGHT 2 - SNAKY 1 - STRAIGHT 2 - SNAKY 3 - OTHER (Specify)
5. LEAF:	
Basal leaf sheath (seedling): 1 = GLABROUS 2 = PUBESCENT	2 Position of flag leaf (at boot stage): 1 - DROOPING 2 - UPRIGHT
1 Waxiness: 1 - ABSENT (Glossy) 2 - SLIGHTLY WAXY 3 - WAXY Leaf sheath is waxy	2 0 MM, WIDTH (First leaf below flag leaf)
1 5 CM. LENGTH (First leaf below flag leaf)	1 Anthocyanin in leaf sheath: 1 = ABSENT 2 = PRESENT
6. HEAD:	a ropor the done
Type: 1 - TWO-ROWED 2 - SIX-ROWED	3 Density: 3 = ERECT (Dense)
Shape: 1 - TAPERING 2 - STRAP 3 - CLAVATE 4 - OTHER (Specify) Strap & Parallel	2 Waxiness: 1 = ABSENT (Glossy) 2 = SLIGHTLY WAXY 3 = WAXY
3 = 1/4 - 1/2 OF HEAD	Rachis (Hair on edge): 1 = LACKING 2 = FEW 3 = COVERED
7. GLUME: 1 - 1/3 OF LEMMA 2 - 1/2 OF LEMMA Length: 3 - MORE THAN 1/2 OF LEMMA	3 Hairs: 1 - NONE 2 - SHORT 3 - LONG
4 Hair covering: 1 - NONE 2 - RESTRICTED TO MIDDLE	3 - CONFINED TO BAND 4 - COMPLETELY COVERED
2 Awns: 1 - LESS THAN EQUAL TO LENGTH OF GLUMES 2 3 - MORE THAN EQUAL TO LENGTH OF GLUMES	- EQUAL TO LENGTH OF GLUMES
Awn Surface: 1 - SMOOTH 2 - SEMISMOOTH 3 - ROUG	н -

		<u> </u>	<u></u>
4 Awn: 3 = SH	YNLESS 2 = AWNLETS ON CENTRAL RO IORT ON CENTRAL ROWS, AWNLETS ON L DNG (longer than spike) 6 = HOODED		RAL ROWS HORT (less than equal to length of spike)
4 Awn Surface: 1	- AWNLESS 2 - SMOOTH 3 - SEMISM	OOTH 4 - ROUGH	
2 Teeth: 1 - ABS	ENT 2 - FEW 3 - NUMEROUS	2 Hair: 1 - ABSE	NT 2-PRESENT
I I Shane of have:	- DEPRESSION 2 - SLIGHT CREASE - TRANSVERSE CREASE	2 Rachilla Hairs:	1 - SHORT 2 - LONG
9. STIGMA:			
Hairs: 1 - FEW	2 - MANY		
10. SEED:	•		
2 Type: 1 - NAK	ED 2 - COVERED	Hairs on Ventral F	urrow: 1 = ABSENT 2 = PRESENT
Z Lenvin:	ORT (8.0 mm.) 2 = SHORT TO MIDLONG DLONG TO LONG (9.0 - 10.5 mm.)		DLONG (8.5 - 9.5 mm.) NG (10.0 mm.)
3 Wrinkling of hull:	1 - NAKED 2 - SLIGHTLY WRINKLED	3 = SEMIWRINKLED	4 - WRINKLED
Aleurone Color:	1 = COLORLESS (White or Yellow) 2 = 8	LUE	enge.
PERCENT AB	ORTIVE	3 2 GMS. PER 10	00 SEEDS
11. DISEASE: (0 = Not	Tested, 1 = Susceptible, 2 = Resistant)		
0 SEPTORIA	1 NET BLOTCH	2 SPOT BLOTCH	2 POWDERY MILDEW
0 LOOSE SMUT	0 BACTERIAL BLIGHT	0 COVERED SMUT	O FALSE LOOSE SMUT
O STEM RUST	2 LEAF RUST	1 SCAB	2 SCALD
O AY	O BSMV	2 BYDV	OTHER (Specify)
12. INSECT: (0 = Not tes	ted, 1 = Susceptible, 2 = Resistant)		
O GREEN BUG	O ENGLISH GRAIN APHID	0 CHINCH BUG	0 ARMYWORM
0 GRASS HOPPERS	1 CERIAL LEAF BETTLE	OTHER (Specify)	
	(O GP O A	0 B 0 C	
HESSIAN FLY RA	CES OD DE	0 F 0 G	
13. CHEMICAL (0 = Not	Tested, 1 = Susceptible, 2 = Resistant)		
О рот	OTHER (Specify)		
14 INDICATE WHICH V	ARIETY MOST CLOSELY RESEMBLES THA	T SUBMITTED:	
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant tillering	Callao	Seed size	Callao
Leef size	Callao	Coleoptile alongation	
Lest size	Callao	Seedling pigmentation	
	Callao		
Leaf carriage		<u> </u>	
REFERENCES: The foll	owing publications may be used as a refer	ence aid for the standar	dization of character descriptions and 🦈 🔻

terms used in this form:

1. Wiebe, G. A., and D. A. Reid, 1961, Classification of Barley Varieties Grown in the United States and Canada in 1958, Technical Bulletin No. 1224, U.S. Dept. of Agriculture.

2. Reid, D. A., and G. A. Wiebe, 1968, Barley: Origin, Botany, Culture, Winter Hardiness, Genetics, Utilization, Pests, Agriculture Handbook No. 338, U.S. Dept. of Agriculture. pp. 61 - 84.

3. Malting Barley Improvement Association, Milwaukee, Wisconsin, 1971, Barley Variety Dictionary.

COLOR: Nickerson's or any recognized color fan may be used to determine color of the described variety. FORM LPGS-470-5 (8-80) (REVERSE)

'PRICE' WINTER FEED BARLEY

18D. EXHIBIT D: ADDITIONAL DESCRIPTION OF PRICE

Since Price has not been tested in comparison with any of the seven cultivars listed in Exhibit C, performance data are presented in Tables 1-15. Price is a high yielding, moderately-early maturing, short awned, six-row winter-feed barley with very good straw strength and test weight. On average, head emergence of Price is 1-2 days later than 'Nomini', 3 days later than Callao, and 2 days earlier than 'Wysor' (Tables 1,8,10,12). The average plant height of Price (36 inches) is 2 inches taller than Callao, 4.5 inches shorter than Wysor and 5.5 inches shorter than Nomini. Straw strength (0=no lodging, 10=completely lodged) of Price (1.4) is excellent in comparison with Nomini (1.8), Wysor (2.0), and Callao (5.9). Average grain yields of Price (114.4 Bu/ac) in Virginia have been similar to those of Nomini and Callao (Table 1). In the Uniform Winter Barley Yield Nurseries (1998-99 and 1999-2000), average grain yields of Price (102 Bu/ac) were similar to those of Nomini, and 3.5 Bu/ac higher than those of Callao (Tables 8, 10). Over all locations (11-12 each year), Price ranked 2nd among 20 entries in 1998, 1st among 18 entries in 1999 and 9th among 25 entries in 2000. Average test weight of Price (50.3 LB/Bu) in Virginia has been only slightly lower (0.7 LB/Bu) than that of Callao, but has been significantly higher than those of Wysor (48.5 LB/Bu) and Nomini (47.5 LB/Bu). In the 1998-99 and 1999-2000 uniform nurseries, average test weight of Price (47.4 LB/Bu) was 0.5 LB/Bu lower than that of Callao, but nearly 3 LB/Bu higher than those of Wysor and Nomini. Price has expressed good winter-hardiness (92% survival) in comparison with Wysor (95%), a very winter-hardy cultivar (Tables 10, 12). In the 1998-2000 Uniform Barley Winter Hardiness Nursery, Price had a mean survival score of 72%, compared with 63% for 'Tennessee Winter', 82% for 'Kentucky 1', and 45% for the winter-tender check 'Trebi' (Tables 13-15).

In adult plant stages, Price is resistant to powdery mildew (*Blumeria graminis* f. sp. *hordei*) and barley yellow dwarf (Tables 2-6, 8, 10). It is moderately resistant to leaf rust (*Puccinia hordei*) and scald (*Rhynchosporium secalis*). It is moderately susceptible to net blotch (*Pyrenophora teres*).

Table 1. Five-year summary of performance of VA96-44-321 in the Virginia Tech Barley Test, 1997 - 2001 harvests.*

Line	Yield (Bu/acre) (23)	Test Weigh (Lb/bı (22)	nt	Date Heade (Mar. 3 (18)	ed 1+)	Heig (In) (16)		Lodgin (0.2-1 (21)	Ō)
VA96-44-321	114.4	50.3	+	20.6		35.6		1.4	
CALLAO	114.3	51.0	+	17.3	-	33.7		5.9	+
NOMINI	114.2	47.5	-	18.8		41.7	+	1.5	· •
STARLING	114.1	47.4	_	22.2	+	40.3	+	2.5	
WYSOR	107.4 -	48.5	-	22.4	+	41.3	+	2.6	
Test Average (n=5)	112.9	49.0		20.3		38.5		2.8	
L.S.D. (0.05)	2.9	0.3		0.4		0.5		0.5	
C.V.	8.4	2.0		6.1		3.8		62.2	

^{*} A plus or minus sign indicates a performance significantly above or below the test average.

The number in parentheses below column headings indicates the number of location-years on which data are based.

[♥] Belgian Lodging Scale = Area x Intensity x 0.2. Area = 1-10, where 1 is barley unaffected and 10 is entire plot affected and Intensity = 1-5, where 1 is barley standing upright and 5 is barley totally flat.

Table 2. Summary of performance of VA96-44-321 in the Virginia Tech Barley Test, 2001 harvest.*

and the second s							Spring	
	Yield (Bu/acre)	Test Weight (Lb/bu)	Date Headed (Mar. 31+)	Height (In)	Lodging v (0.2-10)	Leaf Rust (0-	Freeze Damage 9) •	
Line	(5)	(4)	(4)	(3)	(4)	(1)	(1)	
VA96-44-321	110 +	50.2 -	25 +	31 -	2.1	4 +	4	
CALLAO	104	50.1 -	20 -	30 -	7.7 +	3	5+	
NOMINI	106	47.7 -	23 -	38 ÷	1.8 -	4 +	2 -	
STARLING	103	47.8 -	24	36 +	5.3 +	3	4	
WYSOR	98	48.0 -	25 +	37 +	4.6 +	8 +	3 -	
Test Average (n=41)	101	52.2	24	33	3.1	3	4	
L.S.D. (0.05)	7	0.6	1	1	1.1	1	1	
C.V.	10	1.6	8	3	50.5	23	13	

^{*} A plus or minus sign indicates a performance significantly above or below the test average. The number in parentheses below column headings indicates the number of locations on which data are based.

[♥] Belgian Lodging Scale = Area x Intensity x 0.2. Area = 1-10, where 1 is barley unaffected and 10 is entire plot affected and Intensity = 1-5, where 1 is barley standing upright and 5 is barley totally flat.

[◆] The 0-9 ratings indicate degree to which plant is affected, where 0 = none and 9 = total plant is affected.

Table 3. Summary of performance of VA96-44-321 in the Virginia Tech Barley Test, 2000 harvest*.

	Yield	Test Weight	Date Headed	Height	Lodging&	Leaf Rust	Powdery Mildew
	(Bu/A) (4)	(Lb) (4)	(Mar. 31+) (4)	(In) (3)	(0.2-10) (5)	(2)	0-9) ◊ (2)
VA96-44-321	116	51.4 +	17	37	1.0 -	5	0
CALLAO	115	52.0 +	15 -	35 -	4.4 +	5	ŏ
NOMINI	130+	47.6 -	16 -	43 +	1.0 -	5	0
STARLING	121	47.4 -	20 +	42 +	1.3	4 -	1
WYSOR	112	48.9 -	20 +	43 +	1.8	7 +	Ō
Test Average (n=31)	114	50.6	17	37	1.9	5	1
L.S.D. (0.05)	8	0.7	1	1	0.8	1	
C.V.	10	1.9	5	3			-

A plus or minus sign indicates a performance significantly above or below the test average. The number in parentheses below column headings indicates the number of locations on which data are based.

[♣] Belgian Lodging Scale = Area X Intensity X 0.2. Area = 1-10, where 1 is barley unaffected and 10 is entire plot affected and Intensity=1-5, where 1 is barley standing upright and 5 is barley lying totally flat.

[♦] The 0-9 ratings indicate relative disease intensity where 0=none and 9=total plant infection.

Table 4. Summary of performance of VA96-44-321 in the Virginia Tech Barley Test, 1999 harvest.*

Brand/Variety	Yield (Bu/A)	Test Weight (Lb)	Date Headed (Mar. 31+)	Height	Lodging 4	Powdery Mildew	Leaf Rust	
•	(4)	(LD) (4)	(3)	(In <u>)</u> (3)	(0.2-10) (4)	(0-9 (1)	")◇ (1)	
VA96-44-321	133 +	50.3	22 +	37	0.3 -	1	5	
STARLING	129	47.9 -	24 +	40 +	0.9	1	3 -	
WYSOR	128	48.8 -	24 +	42 +	0.9	1	9 +	
NOMINI	127	47.6 -	21 +	43 +	0.6	1	7 +	
CALLAO	125	51.2 +	19 -	35 -	4.0 +	1	6 +	
Test Average (n=26)	127	50.1	20	27				٠٠.
	1 12/	50.1	20	37	0.9	I	5	
L.S.D. (0.05)	6	0.5	1	1	0.6	. 1	1	

^{*} The number in parentheses below column headings indicates the number of locations on which data are based. A plus or minus sign indicates performance significantly above or below the test average, respectively.

[♣] Belgian Lodging Scale = Area X Intensity X 0.2. Area = 1-10, where 1 is barley is unaffected and

¹⁰ is entire plot affected and Intensity=1-5, where 1 is barley standing upright and 5 is barley lying totally flat.

[♦] The 0-9 ratings indicate relative disease intensity where 0=none and 9=total plant infection.

Table 5. Summary of performance of VA96-44-321 in the Virginia Tech Barley Test, 1998 harvest.*

Brand/Variety	Yield (Bu/A) (5)	Test Weight (Lb) (5)	Date Headed (Mar. 31+) (4)	Height (In) (4)	Lodging (0.2-10)	Scald (0-9)◊ (1)	Leaf Rust (0-9) (1)	Net Blotch (0-9) (1)
VA96-44-321	108 +	47.9 +	18 +	- 36	2.8 -	4	3	5 +
CALLAO	112 +	48.1 +	15 -	33 -	7.6 +	5	3	2 -
NOMINI	106	45.8	14 -	41 +	2.5 -	5	3	1 _
STARLING	109 +	44.8 -	19 +	40 +	3.7	6	2 -	1 - 4 +
WYSOR	96 -	45.4 -	20 +	41 ÷	3.9	5	5 +	3
Test Average (n=40)	103	46.3	17	36	4.3	5	2	3 -
S.D. (0.05)	5	0.8	1	1	1.2	2	1	1

^{*} The number in parentheses below column headings indicates the number of locations on which data are based. A plus or minus sign indicates performance significantly above or below the test average, respectively.

[♣] Belgian Lodging Scale = Area X Intensity X 0.2. Area = 1-10, where 1 is barley is unaffected and 10 is entire plot affected and Intensity=1-5, where 1 is barley standing upright and 5 is barley lying totally flat.

[♦] The 0-9 ratings indicate relative disease intensity where 0=none and 9=total plant infection.

Table 6. Summary of performance of VA96-44-321 in the Virginia Tech Barley Test, 1997 Harvest.*

Brand/Variety	Yield (Bu/A) (5)	Test Weight (Lb) (5)	Date Headed (Mar 31+) (3)	Height (In) (3)	Lodging (0.2-10) 4	Barley Yellow Dwarf Virus (0-9) (1)
VA96-44-321 CALLAO NOMINI STARLING WYSOR	115 122 + 114 120 115	51.9 53.7 + 49.3 - 49.9 - 51.7	23 + 20 - 21 - 26 + 25 +	37 - 36 - 43 + 41 + 43 +	1.2 5.0 + 1.9 1.2 1.6	1 + 0 1 + 0 1 +
Test Average (n=40) L.S.D. (0.05)	114 _. 7	51.8 0.8	22 1	39	1.8 1.1	0

^{*} The number in parentheses below column headings indicates the number of locations on which data are based. A plus or minus sign indicates performance significantly above or below the test average, respectively.

Belgian Lodging Scale = Area X Intensity X 0.2. Area = 1-10, where 1 is barley is unaffected and 10 is entire plot affected and Intensity=1-5, where 1 is barley standing upright and 5 is barley lying totally flat.

[♦] The 0-9 rating indicates relative disease intensity where 0=none and 9=total plant infection.

Stations of the Page 2000 Uniform Winter Barley Yield Nursery			C	1	Les over all sta	nons in the	1999-200	0 Uniform	Winter Barle	y Yield Nu	rserv			
Selection or	Yield	Yield	₹	KĄ.	MD	NC	NE	НО	PA	SC	SC	λL	1 1 1	
variety	Kank	(Pnq)	Griffin	Lexington	Queenstown	Kinston	Lincoln	Wooster	University	Clemson	Florence	.	۷ : ۱	Α Α
Price	6	102.3	109.1	797	81 5	0.00			Park		דווסי בוורב	rrosper	Blacksburg	Warsaw
;						80.9	8/.4	123.2	124.0	113.0	94.9	080	1137	1000
Callao	12	100.1	106.2	9.49	78.9	80.2	78.9	6.68	138.7	112.0	0.300		1.0.1	7.771
Wysor	17	93.0	107.9	74.4	7 7 1	00.0	, ,	į		112.0	102.0	111.9	111.7	123.5
Nomini	v		6	,	-	70.7	67.3	0.//	123.0	94.6	70.9	82.0	116.4	112.6
	J	104.	170.8	95.0	85.7	99.2	105.1	100.2	123.0	110.0	7 00	,	•	
Perkins	23	81.1	85.1	647	0 5 0	ī	i t			0.01	70.4	C.0/	123.0	120.3
i - 				1		C.4.	C./8	91.6	94.0	74.0	69.2	85.7	69.1	,
VA97B-178	9	103.8	113.1	89.3	75.3	91.2	88.1	010	1240	000	! .		1.00	73,3
VA97B-176	2	112.9	1302	75.4		4		7117	0.401	108.0	104.0	104.9	117.1	129.3
		ì	1001	t. C.	92.00	110,8	87.8	109.3	141.0	118.0	116.0	115.5	123.0	130 €
VA97B-388	,	114.0	138.6	83.9	91.3	99.2	976	1367	1710	-			(15):	130.0
Average (N=25)		95.4	1015	977	1 2			130.1	141,0	114.0	118.0	95.2	119.3	133.0
			0.101	4.0	/8/	82.0	88.2	100.3	123.0	94.0	84.0	F 00	0 101	
LSD (0.05)		8.9	24.2*	10.5*	21.2	15.2	18.8*	*8 91	17.0*	7 2 2	; ;	† · oo	6./01	1.6.4
C.V. (%)		5	140	· •	,	,	!		2:	2.0	16.0	13.7	10.6*	10.3*
			14.0	2.0	16,4	9	V V	10.4	d	•			٠	

¹Results of 2 reps. * LSD (0.10)

10.6*

13.7 17.5

16.0 11.0

12.6

15.5

Table 8. Performance of Price versus check varieties in the 1999-2000 Uniform Winter Barley Nursery: Averages

over all stations for indicated characters.

Selection or Variety	Yield Rank	Yield (bu/A)	Test Weight (Ibs/bu)	Heading Date (Julian)	Height (in.)	Lodging (0-9) ¹	Leaf Rust	Powdery Mildew (0-9)
Price	9	102.3	47.0	106	32.8	1.1	2.5	0.5
Callao	12	100.1	47.8	105	32.3	4.4	2.0	0.0
Wysor	17	93.0	46.4	108	35.8	2.5	3.5	0.0
Nomini	5	104.1	46.5	106	37.9	1.8	2.5	0.0
Perkins	23	81.1	48.5	114	36.5	3.8	4.5	1.5
VA97B-178	6	103.8	47.0	107	32.2	2.7	3.0	0.5
VA97B-176	2	112.9	47.0	104	32.6	2.1	2.5	0.5
VA97B-388	1	114.0	47.0	108	34.9	1.3	4.0	0.5
Average (N=25)		95.4	47.5	109	35.5	2.9	3.7	1.3
LSD (0.05)		8.9	1.8	2.0	1.5	1.7	ns²	1.7
C.V. (%)		11.5	4.6	2.2	4.7	51.0	19.8	62.2
No. Locations		12	12	11	10	6	2	2

¹ All 0-9 ratings indicate relative disease/lodging severity: 0 = no disease/lodging present; 9 = total infestation of the plant by disease/completely lodged.

² Not significant.

1 able 9. Kanked yield averages of Price versus check varieties over	yield avera	iges of Pric	e versus c	heck varieties	over all station	S in the 10	71 00 800							
Selection or	Yield	Vield	GA	KY	MD	No.	100 66-92 100 66-92	rorm wint	Mh NC N.	eld Nursery				
Variety	Rank		}			֝֝֝֝֝֝֝֝֝֝֝֝֡֝֟֝	zi Z	HO	PA	\mathbf{sc}	SC	TX	VA	\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \
	THE PARTY OF THE P		Griffin	Lexington	Queenstown	Kinston	Lincoln	Wooster	University		Ā			V.
Price	_	102.1	903	124.4	9 10				Park	Cicinson	Florence	Prosper	Blacksburg	Warsaw
į		•		1747	17/.8	97.0	67.5	152.8	1150	702	0.53			
Callao	9	97.1	99.4	106.5	1107	0 5 0	9			0.0	51.8	56.4	123.0	149.0
Wysor	10	0.0	9	1		0.00	07.70	130.0	115.0	58.9	54.7	0.69	126.0	147.0
	27	75.4	7.7/	118.6	119.2	81.0	82.8	6 96	108.0	7 36		i) : :	2
Nomini	m	9.66	9.09	118.8	100 0	ç	:		100,0	0.07	45.0	70.7	117.0	132.0
Dorling	Ţ	6			120.0	98.0	73.1	137.3	111.0	93.7	41.5	889	122.0	4
T CI WIIIS	<u>`</u> 1	80.2	73.6	103,4	86.5	81.0	. 1 09	00 7	100				0.771	147.0
VA97B-178	7	100.1	0 06	1002					102.0	20.0	35.0	60.5	101.0	109.0
100				107.7	11/11	90.0	45.7	137.8	130.0	69.7	40.9	80.2	1250	\ \ \
Average (N=18)		92.9	77.1	114.7	111.4	84.0	643	124.5	105.0	9		7:00	0.001	156.0
LSD (0.05)		11.2	22.2	10.7	-)	77.7	102.0	0.80	43.9	64.4	123.0	133.0
		!	7:1	10.1	18.0	12.0	21.1	20.2	18.0	6.7	~	90		•
C.Y. (%)		14.9	13.8	5.4	10.0	9.8	23.8	ă O			;	0.7	0.11	13.0

Table 10. Performance of Price versus check varieties in the 1998-99 Uniform Winter Barley Nursery: Averages over all stations for indicated characters.

Selection or Variety	Yield Rank	Yield (bu/A)	Test Weight (lbs/bu)	Heading Date (Julian)	Height (in.)	Lodging (%)	Winter Survival (%)	Leaf Rust	BYDV (0-9)
Price	. 1.	102.1	47.8	112	32.1	5.1	91.2	5.3	1.9
Callao	6	97.1	47.9	110	30.9	37.2	93.5	5.3	1.1
Wysor	10	93.4	46.0	114	35.9	14.6	94.4	7.3	1.5
Nomini	3	99.6	46.1	113	36.5	17.2	92.9	6.0	1.6
Perkins	17	80.2	48.7	119	36.0	13.5	99.5	4.0	2.5
VA97B-178	2	100.1	47.7	111	31.2	5.4	84.3	4.3	1.0
Average (N=18)		92.9	46.9	114	33.9	13.8	89.0	5.7	1.5
LSD (0.05)		11.2	3.0	9.4	1.8	14.7	21.1	1.9	ns ²
C.V. (%)		14.9	8.1	2.1	6.4	93.0	16.7	19.8	
No. Locations		12	12	10	12	6	4	3	2

¹ All 0-9 ratings indicate relative disease severity: 0 = no disease present; 9 = total infestation of the plant by disease.

² Not significant.

Selection or Viola Viv. GA KV CV	Viola	*7.	G.A.	7.7		אוז חוז חוז כווכ	27/-78 Un	Itorm Wint	er Barley Yie	old Nursery			
Variety	י וכנות	r Jeld	;	4	QIM	NC	Z	НО	PΑ	50			
(Table)	Kank	(pn/A)	Griffin	Kank (bu/A) Griffin Lexington	Queenstown	Kinston	Inoni	W	University	ر د) N	VA	ΛA
Price	2	88.7	100.6	\$7.0				w ooster	Park	Clemson	Florence	Blacksburg	Warsaw
Wysor	8	77.5	41.1		5.0%	53.2	88.9	122.4	0.66	84.6	48.0	1050	136.0
	2	C.7	4.1.1	8.67	83.5	38.9	79.7	92.5	03.0		;		120.0
Militon	20	68.7	61.7	26.1	1 92	, 40	i	i i	2:0	/'C/	20.8	0.66	113.0
Average (N=20)		82.2	747	17.4		55.3	72.7	80.4	104.0	59.6	39.4	0.06	110.0
130 W 057		0		†. †	84.8	51.3	6'98	111.9	102.7	77.9	48.0	0 00	
(cara)		y. 8.	8.4		12.6	11.0	16.1	20.0	15.0		ì	70.0	119.0
C. V. %	-	14.2	•		ć	,		: } }	0.01	11.0	8.0	0.6	14.0
					9.0	14.5		100					

Table 12. Performance of Price versus check varieties in the 1997-98 Uniform Winter Barley Nursery: Averages over all stations for indicated characters.

Selection or Variety	Yield Rank	Yield (bu/A)	Test Weight (lbs/bu)	Heading Date (Julian)	Height (in.)	Lodging (%)	Winter Survival (%)
Price	2	88.7	46.0	114.1	31.5	21.9	92.3
Wysor	18	72.5	42.6	114.7	37.6	35.8	96.3
Milton	20	68.7	45.1	117.0	34.1	50.5	89.6
Average (N=20)		82.2	44.3	115.0	35.8	35.4	93.6
LSD (0.05)		9.8	1.4	2.0	1.5	19.4	6.6
C.V. (%		14.2	3.8	2.0	5.3	55.4	5.0
No. Locations		11	11 -	10	12	. 6	3

Table 13. Percent winter survival of Price versus check varieties at the various stations in the 1999-2000

Uniform Barley Winter Hardiness Nursery.

Selection or Variety	Rank According	_	Canada	Turkey	NE	NE	TX	Rank According	Means Across	KS
	to Mean	Locations	Nairn	Eksehir	Lincoln	Mead	Dallas	to Mean	Locations ²	Manhattan ³
Price	8	88.0	90.0	50.0	100.0	100.0	100.0	3	89.1	100.0
Tenn. Winter	22	81.0	77.5	37.5	95.0	95.0	100.0	23	73.6	0.0
Trebi	29	63.0	12.5	7.5	100.0	95.0	100.0	29	57.3	0.0
Kearney	19	83.0	80.0	35.0	100.0	100.0	100.0	13	84.5	100.0
Kenosha (wheat)	9	87.8	99.0	55.0	100.0	85.0	100.0	4	88.9	100.0
Dicktoo	12	86.5	95.0	52.5	100.0	85.0	100.0	7	87.7	100.0
Kentucky 1	15	85.0	92.5	42.5	90.0	100.0	100.0	11	86.4	100.0
VA97B-178	6	90.0	80.0	70.0	100.0	100.0	100.0	2	90.9	100.0
VA97B-176	5	90.3	89.0	62.5	100.0	100.0	100.0	10	86.6	50.0
VA97B-388	1	93.5	97.5	70.0	100.0	100.0	100.0	12	85.5	5.0
Average (N=29)		83.6	76.3	46.3	99.1	96.9	99.1		80.0	43.5
LSD (0.05)		18.6	14.3	23.2	ns ⁴	6.6	0.9			
C.V. (%)		11.7	14.5	38.3		5.3	0.7			

¹ Means obtained by MSTAT analysis excluding data from Manhattan location.

² Unweighted means calculated for the locations including Manhattan.

³ Data from single replication of one meter row plots.

⁴ Not significant.

Table 14. Percent winter survival of Price vers

Selection or	Kalik	Means	δ	=	07	GA ID VS		13111 II (2)	rial diness p	lursery.
Variety	According	Across	; }	3	2	MD	NE	NE	PA	SD
	to Mean	Locations	Griffin	Aherdean	E Lossia				University	
Price	13	62.9	08	Tan Isaac	riays	Keedysville	Lincoln	Mead	Park	Brookings
Tenn. Winter		7	00	87	100	80	09	45	100	5
	,	45.1	80	49	83	70	10	v		
Trebi	. 56	22.6	70	14	75	٥	?		3	0
Kearney	7	90.0	80	86	100	o {	-	0	. 65	0
Kenosha (wheat)	. —	92'6	8	2 001		93	100	06	100	09
Dicktoo	ij	0 98	} :	3	MI t	56	100	100	100	82
Kontuolee 1) \	600	6 3	%	100	95	100	85	100	80
rentucky 1	9	83.8	70	83	86	88	100		ć	
VA97B-178	19	9.99	75	28	001) (3	C ₁	86	09
Average (N=26)		64.0			3	90	40	20	100	10
6-1-1-0		04.y	/3	28	95	78	56	38	70	
LSD (0.05)		19.0	ns	50	5	12	70) i	0.6	97
C.V. (%)		216	2	Ç	} '	71	6	72	=	19
		71.0	71	747	∞ ·	∞	42	33	ĸ	35

¹ Not significant.

Table 15. Percent winter survival of Price versus check varieties at the various stations in the 1997-98 Uniform Barley Winter

Hardiness Nursery.

Trandiness radisery								
Selection or Variety	Rank According	Means Across	Canada	NE	NE	PA	SD	WI
v ariety	to Mean	Locations	Nairn	Lincoln	Mead	University Park	Brookings	Madison
Price	19	64.4	94.0	95.0	35.0	100.0	20.0	42.5
Tenn. Winter	21	62.8	100.0	95.0	25.0	97.0	20.0	40.0
Trebi	23	48.9	100.0	80.0	35.0	60.0	10.0	8.5
Kearney	. 11	72.8	100.0	100.0	55.0	100.0	estd=29.5	52.5
Kenosha (wheat)	1	92.0	100.0	100.0	100.0	100.0	estd=53.0	99.0
Dicktoo	6	75.4	100.0	95.0	40.0	100.0	60.0	57.5
Kentucky 1	4	76.7	100.0	100.0	60.0	100.0	45.0	55.0
Average (N=23)		71.7	99.5	97.8	52.2	97.9	30.0	50.2
LSD (0.05)		14.2	2.2	ns¹	31.1	6.7	ns	10.2
C.V. (%)		17.4	1.1	4.7	28.7	3.3	85.0	9.8

¹ Not significant.

REPRODUCE LOCALLY, Include form number and edition date on a	III reproductions.	FORM APPROVED - OMB No. 0581-0055
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E	Application is required in order to det certificate is to be issued (7 U.S.C. 2 confidential until the certificate is issued.	421). The information is held
STATEMENT OF THE BASIS OF OWNERSHIP 1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
	OR EXPERIMENTAL NUMBER	
Virginia Tech Intellectual Properties, Inc.	VA96-44-321	Price
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)
1872 Pratt Drive, Suite 1625 Blacksburg, VA 24060	540-951-9374	540-951-5292
	7. PVPO NUMBER	300132
8. Does the applicant own all rights to the variety? Mark an "X" in the	ne appropriate block. If no, please expl a	ain. YES NO
9. Is the applicant (individual or company) a U.S. national or a U.S.	based company? If no, give name of o	country. YES NO
10. Is the applicant the original owner?	NO If no, please answer one	of the following:
a. If the original rights to variety were owned by individual(s), is YES YES	(are) the original owner(s) a U.S. Nation NO If no, give name of count	
b. If the original rights to variety were owned by a company(ies	s), is (are) the original owner(s) a U.S. ba	
11. Additional explanation on ownership (If needed, use the reverse	e for extra space):	
Original owner Virginia Polytechnic Institute and State Tech Intellectual Properties, Inc. (See Attached)	University assigned its ownersh	ip to current owner Virginia
PLEASE NOTE:		
Plant variety protection can only be afforded to the owners (not lice		
 If the rights to the variety are owned by the original breeder, that national of a country which affords similar protection to nationals 	person must be a U.S. national, national of the U.S. for the same genus and spec	l of a UPOV member country, or cies.
If the rights to the variety are owned by the company which emple nationals of a UPOV member country, or owned by nationals of a genus and species.	oyed the original breeder(s), the compar a country which affords similar protection	ny must be U.S. based, owned by to nationals of the U.S. for the same
3. If the applicant is an owner who is not the original owner, both the	e original owner and the applicant must	meet one of the above criteria.
The original breeder/owner may be the individual or company who out for definitions.	directed the final breeding. See Section	41(a)(2) of the Plant Variety Protection
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponso control number. The valid OMB control number for this information collection is 0581-005 including the time for reviewing the instructions, searching existing data sources, gathering	The time required to complete this information coll	ection is estimated to average 0.1 nour per response,

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY (hereinafter referred to as the "UNIVERSITY"), assigns to VIRGINIA TECH INTELLECTUAL PROPERTIES, INC. (hereinafter referred to as "VTIP") all rights, title and interest in and to the GERMPLASMS listed below as held by the UNIVERSITY:

· · · · · · · · · · · · · · · · · · ·
Price/VA96-44-321 Barley
VA98W-593 Wheat
VA97W-469 Wheat
McCormick/VA98W-591

The UNIVERSITY, by its authorized agents, agrees that it will execute all necessary assignments as requested by VTIP, to facilitate the filing of patent applications and/or copyright registrations. It will render any reasonable assistance requested to aid in preparation of such applications and/or registrations.

The UNIVERSITY shall retain the right to make use of the GERMPLASM for internal research and other non-commercial purposes without cost to the UNIVERSITY.

All royalties, rents, payments, or any cash receipts from the sale, assignment, transfer, licensing or use of the GERMPLASM shall be the property of VTIP and shall be distributed according to the provisions of the Virginia Agricultural Experiment Station (VAES) Plant Germplasm Release Policy (PGRP).

Prior to the execution of this Assignment, the UNIVERSITY has not granted the right of license to make, use, or sell said GERMPLASM to anyone except to VTIP, nor has it otherwise encumbered its rights, title and interest in said GERMPLASM, and it will not execute any instrument in conflict with this Assignment.

> VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY

BY

MINNIS E. RIDENOUR Chief Operating Officer

STATE OF VIRGINIA

COUNTY OF MONTGOMERY, to-wit:

The f	oregoing instrument was acknowledged before me this 18th day of
APRIL	, 2002, by MINNIS E. RIDENOUR CHIEF OPERATION
of Virginia P	olytechnic Institute and State University, on behalf of said University.
	Notary Public
	My commission aminos 12/31/04